



# Hydrogeological researches for still waters in Codru Moma and Pădurea Craiului Mountains (Apuseni Mountains, Romania)

Iancu Orășeanu  
S.C. *Prospecțiuni S.A.*

RESUME. Afin d'identifier des sources d'eau plate dans les montagnes Codru Moma et Pădurea Craiului, on a soumis un nombre de 6 sources et la sonde S5 Moneasa a une recherche hydrogeologique complexe. Les sources ont été mises en évidence comme probablement d'eau plate par des investigations antérieures, au caractère régional.

La recherche hydrogeologique complexe a consisté en établir les conditions hydrogeologiques dans lesquelles apparaissent les sources, l'observation à la source de la variation des paramètres physico-chimiques pendant au moins une année et établir la variation dans le temps de la composition chimique et de la charge bactériologique, sur des échantillons d'eau mis en bouteille et stockés dans des conditions de laboratoire. On a également analysé le contenu en éléments toxiques et en pesticides des eaux, ainsi que leur radioactivité.

Les sources étudiées ne sont pas aménagées, raison pour laquelle certaines d'entre elles présentent temporairement des charges bactériologiques élevées, dues à une contamination locale, à la source (Feredeu, Izvorul Mare al Tărcăiței, Țucrești).

Les eaux sont de type Ca Mg HCO<sub>3</sub> ou MgCa HCO<sub>3</sub>, avec des minéralisations faibles (moins de 500 mg/l résidu sec), sans éléments toxiques et non-radioactifs. Le contenu en pesticides dépasse chez certaines sources de 2 fois la concentration maximale admise par les normes de potabilité. Les eaux ont une bonne stabilité chimique, à l'exception de la source Țucrești, laquelle est sursaturée par rapport au calcite, ce-dernière présentant avec le temps de tendances à précipiter.

Le débit des sources est assez grand pour assurer une exploitation économique et présente une variabilité faible avec le temps, sous ce dernier aspect se distinguant les sources Feredeu et Rășchirata.

La sonde S5 Moneasa donne de l'eau plate.

## 1. INTRODUCTION

The regional hydrogeological investigations performed in the Codru Moma Mountains, outlined the occurrence of good quality groundwater sources, some of which can be bottled as table water.

These sources were subject of a complex investigation program, conducted by S.C. *Prospecțiuni S.A.*, over a two period of time, 1991-1992 and 1994-1996, that resulted in:

- outlining the physico-chemical parameters of the sources, by means of systematic discharge, temperature, pH, electrical conductivity, dissolved oxygen, alkalinity and redox potential measurements;
- establishing the long term evolution of the chemical composition (especially labile elements and parameters: HCO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>-2</sup>, Ca<sup>++</sup>, Mg<sup>++</sup>, dissolved oxygen, total hardness, residue on evaporation at 105<sup>0</sup>C) and of the bacteriological content, both at the source and for samples bottled and stored in laboratory conditions;
- outlining the toxic elements and pesticides content, and the global  $\alpha$  and  $\beta$  radioactivity.

In the framework of the conducted study, the laboratories of *Prospecțiuni S.A.* have performed the general chemical analyses and the toxic elements contents measurements. The bacteriological analyses have been performed at the Center for Prophylactic Medicine in Beiuș, under the leadership of dr. D. Mocuța, and partly at the Food Chemistry Institute in Bucharest.

For determining the pesticides content, the water samples have been analysed in the Central Laboratory for the Control of Animal and Vegetal Origin Products in Bucharest, while

the total  $\alpha$  and  $\beta$  radioactivity has been performed by the laboratories of the Institute of Physics and Nuclear Engineering in Măgurele and in the laboratories of the Rare Metals Authority in Măgurele.

## 2. INVESTIGATED SOURCES

Codru Moma Mountains are situated in the western part of the Apuseni Mountains and they occur as a massive body which extends between the Neogene basins of Beiuş and Zarand. Moneasa and Crişul Văratec streams separate two distinct physiographic units: Codru Mountains - to the north, and Moma Mountains - to the south.

Codru Moma Mountains prevalently consist of Permian-Mesozoic sedimentary deposits. In the geologic constitution of their northern half are included Triassic and Early Jurassic limestones and dolomites, which form a continuous N-S striking strip. In the Moma Mountains, Triassic carbonate deposits form the lithologic background of the Vaşcău karst plateau (fig. 1).

The complex hydrogeological investigations program has included 6 springs (table 1). The program was also aimed at outlining the quality of water discharged by the well S5 in Moneasa spa.

Table 1. Characteristics of the sources.

No	Source	T °C	Q l/s	TDS mg/l	RE mg/l	TH °d	S.I.		Pest. ppb	$\alpha+\beta$ mBq	Bac	
							lms	dol			a	b
1	Feredeu (Huta, Finiş)	11,2-13,0	55,0	518,1	316,2	16,03	0,038	0,086	0,0005		+	-
2	Izv. Mare al Tărcăitei	10,0-10,6	70,0	687,7	285,2	15,15			0,0008	< 42	-	+
3	Well S5 Moneasa	14,2-16,4	4,9	302,0	151,8	6,7	-0,181	-0,463	0,0009	81	+	+
4	Răşchirata (Dezna)	8,2-8,8	35,9	639,9	267,5	13,8	0,164	0,004	0,0008	< 60	+	+
5	Valea Seacă (Dezna)	8,1-8,6	63,7	558,3	259,7	13,7	0,040	-0,212	0,0007	< 34	+	+
6	Țucreşti (Vaşcău)	14,4-15,4	5,7	566,7	279,8	14,55	0,577	0,498	0,0007	55	-	+
7	Pancului (Remeţi)	9,5-10,2	29,0	514,6	332,1	15,2			0,0010		+	+

Note: T- temperature; Q-mean annual discharge; RE-residue on evaporation at 105°C; TH-total hardness; S.I.-saturation index (lms-limestones; dol-dolomites); Pest-pesticides (sum of  $\alpha$  HCH,  $\beta$  HCH and lindan);  $\alpha+\beta$  - global radioactivity; Bact.-Result of bacteriological analysis: a- seasonal analysis; b-analysis of bottled and stored samples; (+)-bacteriological content less than maximum accepted contents for drinking water by STAS 1342-91 (m.a.c.); (-)-bacteriological content higher than m.a.c., at least at one sample.

The following general assertions are valid for the investigated springs:

- the springs are either not tapped at all, or tapped in a rudimentary manner, which results in a local bacteriological contamination, mainly ensuing to runoff water inflow. Due to this fact, in some cases concentrations do not comply with drinking water requirements;

- all studied spring waters from Codru Moma Mountains are  $\text{CaMgHCO}_3$  or  $\text{MgCaHCO}_3$  type, with low mineralization (fig. 2);

- the water samples collected from the springs displayed a good chemical stability over the one year monitoring period; The water of Țucreşti spring is supersaturated with respect to calcite, being hence prone to carbonate scaling.

- the saturation indexes, computed by Netpath program, indicate that the water of most karst springs is saturated, to a larger or smaller extent, with respect to both calcite and dolomite. The Țucreşti warm springs is supersaturated with respect to calcite and dolomite. Feredeu, Răşchirata and Valea Seacă springs have saturation indexes very close to the equilibrium;




Fig. 1. Hydrogeological map of Codru Moma Mountains.

Legend. 1-Pannonian-Quaternary deposits (argillaceous shales, marls, sands, gravels; a-colluvium); 2-Neogene magmatites (agglomerates, andesites); 3-Carbonate mesozoic series (limestones, dolomites); 4-Mesozoic flysch-like series; 5 Prevalent detritic Permo-Mesozoic deposits (sandstones, conglomerates and less frequently argillaceous shales and rhyolites) ; 6-Geological boundary; 7-Boundary of colluvium; 8-Fault; 9-Overthrust front; 10-Course of perennial stream; 11-Course of temporary stream; 12-Boundary of endoreic areas; 13-Swallet; 14-Losses in flow along the riverbed; 1-Inflow cave; 16-Mean annual discharge of springs (l/s): a-less than 10; b - 11-50; c - 51-500; 17-Outflow cave; 18-Well with thermal water ( $t > 20^{\circ}\text{C}$ ); 19-Group of springs and wells with subthermal ( $10 < t \leq 20^{\circ}\text{C}$ ) and thermal waters; 20-Selected gas leaks associated with subthermal springs; 21-Ebb and flow spring; 22-Proved groundwater flow direction; 23-Presumed groundwater flow direction.

Key of the numbers on the map: 2-Feredeu spring; 4-Dosul Varului swallet; 5-Grota Ursului spring; 8-Izvorul Mare al Tărcăiței spring; 10-Țucrești spring; 16-Rășchirata spring; 17-Valea Seacă spring.

- the toxic elements content, analysed by means of the ICP equipment in "Prospecțiuni" S.A. laboratories, indicate Cr, Cu, Mn, Ni, Pb, and Zn contents below the maximum allowed concentrations;
- for some sources, the pesticides content exceeds up to 2 times the accepted concentration (0.0005 ppm);
- the waters are not radioactive;
- the gas outflows from Feredeou and Țucrești springs have a composition similar to that of the atmospheric gas;
- the main aquifers in Pădurea Craiului Mountains occur within carbonate rocks having undergone intense karst processes and which are deprived of covering deposits with either good filtering properties, or impervious. They are extremely vulnerable to pollution factors and do not include good quality waters (Orășeanu, 1994).




Fig. 2. Piper diagram with displays of chemical composition of studied springs from Codru Moma and Pădurea Craiului Mountains. (key of the numbers in table no. 1)

## 2. 1. Feredeou spring (figures 1, 2 and table 1)

Feredeou spring is located in the upper catchment area of Finiș stream, on the right side of Bălăteasa tributary, in the area that the locals call Huta (fig. 1, no. 2). Access to the spring became difficult after the Finiș-Huta-Brătcoia narrow-gauge railway was dismantled.

The spring water upflows from between the alluvia which hide the contact between the Carnian-Norian upper dolomites and the Rhaetian marls. The spring recharge area is located further to the south, in the Brătcoia closed basin: the water sinking through Dosu Varului swallet (fig. 1, no. 4) is forced northward in a deep, flooded circulation beneath the marls stack, as tracer tests performed by I. Orășeanu and E. Gașpar in 1977 and 1987 (Orășeanu, 1987; Gașpar & Orășeanu, 1987) have proven. As a result of this deep circulation the spring water is permanently very clear, has a slightly elevated temperature (11.2-13°C), and displays strong gas outflows.

The average Feredeou spring discharge is 55 l/s, with very small fluctuations, ranging between 45 - 70 l/s. Since the spring is not tapped and is invaded by perennial aquatic vegetation, the bacteriological content displays large fluctuations as a function of season.

## 2. 2. Well S5 Moneasa (figures 1, 2, 3, 4 and table 1)

Moneasa resort is located in the central part of Codru Moma Mountains, in an area that underwent strong tectonic displacements, which resulted in an overthrusts structure (fig. 3). The reputation of the resort is due to the thermal water occurrences (both natural springs and drilled wells), with temperatures ranging between 24-32.8°C (Orășeanu, 1987).




Fig 3. Hydrogeological map of Moneasa spa. (Geological data after Bleahu, 1965 & Bleahu et al., 1979, 1984)

Legend. 1-Triassic limestones and dolomites (Id-Ladinian black limestones with cherts; cr-Carnian white dolomites; J<sub>1</sub>-Lower Jurassic red limestones); 2-Quaternary alluvium; 3-Norian argillaceous shales with sandstones and limestones; 4-Permian diabases, sandstones and shales; 5-Course of perennial stream; 6-Geological boundary of quaternary deposits; 7-Fault; 8-Overthrust front; 9-Outflow cave; 10-Thermal spring; 11-Well with thermal water ( $t > 20^{\circ}\text{C}$ ); 12-Well with subthermal water ( $10 < t \leq 20^{\circ}\text{C}$ ).

The karst reservoir has a monoclin, eastward dipping structure, with Werfenian quartzite sandstones at the bottom and flysch type Norian deposits at the top. In the south it is overthrust by the Permian deposits of the Moma nappe (schists, sandstones, basalts).

The thermal water accumulation is the result of mixing between warm water upflows along fractures and cold water recharge from the north. The strong tectonic disturbances underwent by that area and the heterogeneity of its karst are mirrored by the lack of uniformity of the temperature of the water discharged by the wells and by the contrasting TDS contents, with specific time evolutions. The unimodal and narrow distributions are characteristic for thermal sources, the plurimodal and widely ones, for cold waters (Grotă Ursului spring) and, in a little manner, of well S5 (fig. 4).

Well S5 (well no.4666) traversed a succession including Permian schists, sandstones and basaltes belonging to the Moma Nappe (0-275m) and then penetrated Triassic karstified limestones (275-340m) belonging to the Finiș Nappe. The water intercepted in Triassic limestones boast an artesian discharge of 4,6 l/s of water with a average temperature of 16°C.

## 2. 3. Izvorul Mare al Tărcăiței (figures 1, 2, 5 and table 1)

Izvorul Mare al Tărcăiței is located some 13 km south of Beiuș town, on the right side of Tărcăița stream, 2 km upstream the homonym village (fig. 1, no. 8). It occurs as an upflow from alluvial deposits, in close neighbourhood of the northern slopes of Clăptescu

summit, where mainly Anisian dolomites outcrop.

The flood pulses in the spring of 1994 entrenched the streambed of Tărcăița next to the spring by almost 1 m. As a result, the spring discharges entirely into the stream, its former outlet being currently dry.




Fig. 4. Distributions of hydrochemical characters of waters in Moneasa spa.

The spring water, supplied by the karst area existing to the south, undergoes seasonal bacteriological contamination as a result of the cattle breeding performed in that area (fig. 3).




Fig. 5. Displays of the total number of bacteria (N.t.b.) and of the number of coliform bacteria (N.b.c.) in Izvorul Mare al Tărcăiței spring in 1991-1992 time interval.

#### 2.4. Rășchirata spring (figures 1, 2, 6, 7 and table 1)

Carbonate deposits in Vașcău plateau host major water accumulations, that discharge at the plateau border through a few springs displaying high flow rates even in drought periods (Orășeanu, 1985). Two of those sources, Rășchirata spring and Valea Seacă spring, are located at the south-western extremity of the plateau, in the upper catchment area of Zugău stream, which in the Dezna village joins Moneasa stream. A third source monitored in the framework of the program aimed at identifying new still water sources was the spring Țucrești, in the neighborhood of Vașcau town.

Rășchirata spring is located 11 km upstream Dezna village, on the left side of Căptălanul stream, a tributary of Zugău stream, in close neighbourhood of an old iron ore melting furnace and of a forestry hut (fig. 1, no. 16). The spring upflows from the scree slope, at the bottom of a wall where a few meters thick beds of Anisian dolomites outcrop.




Fig. 6. Displays of the total number of bacteria (N.t.b.) and of the number of coliform bacteria (N.b.c.) in Rășchirata spring in 1991-1992 time interval




Fig. 7. Displays of the total number of bacteria (N.t.b.) and of the number of coliform bacteria (N.b.c.) in water sampled from Rășchirata spring at 15.VI.1992, stored and analysed

## 2. 5. Valea Seacă spring (figures 1, 2, 8 and table 1)

Valea Seacă spring is located 1 km upstream Rășchirata spring, on the left side of Valea Seacă, at its confluence with Căptălanul stream (fig 1, no. 17). The water emerges through a couple of closely situated cracks, enlarged by the water. In 1987 the spring has been tapped by the mining exploration company IPEG Deva, the working site Dezna, for the supply of the iron ore washing pilot plant located upstream, at the site called La Grajduri. The water intake is currently dismantled.

In order to delineate the supply area of those two sources, fluorescein tracer was injected on 25 July 1994 in the Scărița swallet (4 l/s flow rate), the tracer being recovered in the Valea Seacă spring (located 1.05 km away and 45 m below) after about 11 hours. The same day, In EDTA tracer was injected in the diffuse total sinking section of Căptălanul stream, located upstream its junction with Valea Seacă stream. 5 hours after its injection the tracer was identified in the Valea Seacă spring as well, 0.6 km away and 25 m below the sinking area. The tracing experiments have been conducted in co-operation with geologist engineer E. Căpraru, and the analysis of the In EDTA tracer concentrations has been performed in the IFIN laboratories in Bucharest.




Fig. 8. Displays of the total number of bacteria (N.t.b.) and of the number of coliform bacteria (N.b.c.) in Valea Seacă spring during 1991-1992.



## 2. 6. Țucrești spring (figures 1, 2, 9 and table 1)

In the western part of Vașcău town, four hypothermal springs (Sfârâșele, Rengle, Racova and Țucrești) emerge from the Late Triassic limestone of the plateau and from the Crișul Negru and Boiu streams alluvia. Their temperatures range between 14.5 and 17.2°C, their cumulated discharge amounts to about 15 l/s, and they display violent gas outflows. Gas outflows display a series of cold springs as well. The springs emerge along the NW-SE striking fractures system, that controlled the progressive Beiuș basin subsidence toward north-east.

The chemical composition of the sub-thermal water at Vașcau is CaMgHCO<sub>3</sub> facies. In some instances the water displays iodide or even bromide character, as a result of reactions with the deposits filling the Beiuș basin. The same chemical character is also met in the case of other sub-thermal outlets in the basin (Ceica, Răbăgani, etc.).

Țucrești spring is located in the western part of Vascau town, at the bottom of Dealul Viilor (fig. 1, no. 10). The spring upflows from the Crișul Negru flood plain alluvia and is permanently invaded by grassy aquatic vegetation.




Fig. 9. Displays of the total number of bacteria (N.t.b.) and of the number of coliform bacteria (N.b.c.) in Țucrești spring during 1991-1992.

## 2. 7. Pancului spring (figures 2, 10, 11, 12 and table 1)

In the eastern part of Pădurea Craiului Mountains is situated the Remeți graben, with Mesozoic limestones and sandstones bordered by crystalline schists and igneous rocks (fig. 10).

Pancului spring is situated within Remeți village, on the right side of the Iad stream, at 450 m a.s.l. (fig. 10, b). The source emerges at the bottom of a steep and grass covered slope debris, that continues with a limestone scarp. To the south and east of Pancului spring carbonate deposits form a synclinal structure, with the Anisian limestone being overlain by Hettangian - Early Sinemurian quartzite sandstones and shales, and further on, in the structure axis, by Late Sinemurian - Carixian limestone (Elena Popa, 1981).

Detritic and clayey Early Jurassic deposits are virtually impervious and force the water which infiltrates along the southern border of the graben, in the Selhișului stream catchment area, into a deep circulation, to re-emerge via Pancului spring. The spring enjoy gas outflows with a composition similar to that of the atmospheric gas (Oășanu, 1991).

T. Rusu (1988) in his comprehensive study concerning the karst in Pădurea Craiului Mountains suggests that Pancului spring might probably drain the water of Selhișului stream. The spring water is clear and only seldom, during heavy rainfall and snow melt periods it has a slightly opalescent colour. Figure 11 displays the total number of bacteria and total coliform bacteria contents of some springs, illustrating a wide range of contents. In the same periods the bacteriological content increase in a moderate way (fig. 12).





Fig. 10. Hydrogeological map of Remeți area. Geological data after Elena Popa (1981).

Legend: 1-Course of perennial stream; 2-Course of temporary stream; 3- Geological boundary; 4-Fault; 5-Subterranean capture; 6- Proved groundwater flow direction; 7- Presumed groundwater flow direction; 8- Alluvium; 9-Mesozoic limestones; 10-Lower jurasic quartzitic sandstones and limestones; 11-Permo-Werfenian sandstones and conglomerates; 12-Rhiolites; 13- Dacites; 14-Crystalline schists.

Key of the numbers on the map: a- Tăul fără Fund spring; b-Pancului spring; c-Spring of Lunca Pizlii; d- Davelii spring; e-Strvinosu spring.



LEGENDA





Fig. 11. Displays of the total number of bacteria (N.t.b.) and of the number of coliform bacteria (N.b.c.) in some springs water Remeți graben.




Fig. 12. Displays of the total number of bacteria (N.t.b.) and of the number of coliform bacteria (N.b.c.) in Pancului spring in 1991-1992 time interval

## CONCLUSIONS

The regional hydrogeological studies performed in Codru Moma and Pădurea Craiului mountains have outlined the occurrence of many karst outlets, among which 6 springs and well S5 from Moneasa have been selected for complex investigations, aimed at the identification of new still water sources.

The investigated outlets are not tapped, and as a consequence some of them temporarily display high bacteriological loads. We assume the phenomenon to be the result of local contamination.

The water discharged by well S5 Moneasa, Rășchirata and Feredeiu springs display a remarkable constancy of the physico-chemical characteristics, circumstance that recommends them as still water sources. Similar properties displays the spring in Valea Seacă as well, its water being however slightly opalescent over short time intervals, during the very high discharge periods. In order to be ascribed to the same category, Pancului spring in Remți requires the hydrologic relationship with the swallets of Selhisului stream to be elucidated.

The water discharged by Izvorul Mare al Tărcăiței is of good quality, unfortunately its sub-aerial outlet has disappeared ensuing to the interception by Tărcăița stream, whose streambed underwent a significant entrenchment during the high discharge period of 1994. In order to be included in the category of the still water sources, the spring must be tapped and the specific analyses must be repeated.

The water of Tucrești spring is supersaturated with respect to calcite, being hence prone to carbonate scaling.

## REFERENCES

- Bleahu, M. (1965) *Harta geologică a RSR, scara 1:100.000, foaia Moneasa*. IGG, București.
- Bleahu, M., Panin Stefana, Camelia Tomescu, Stefan, A., Istrate, Gh., Stefanescu, M. (1979) *Harta geologică a RSR, scara 1:50.000, foaia Vașcau*. IGG București.
- Gășpar, E., Orășeanu, I. (1987) *Natural and artificial tracers in the study of the hydrodynamics of the karst*. Theoretical and Applied Karstology 3, 31-107, București.
- Orășeanu, I. (1985) *Considerations on the hydrogeology of Vașcău Plateau (Codru Moma Mountains)*. Theoretical and Applied Karstology 2, 199-209, București.
- Orășeanu, I. (1987) *Hydrogeological study of Moneasa area (Codru Moma Mountains)*. Theoretical and Applied Karstology 3, 175-199, București.
- Orășeanu, I. (1994) *Considerations regarding the karstic aquifer pollution in Romania, Especially Apuseni Mountains refferings*. Proceedings of the International Hydrogeological Symposium "Impact of industrial activities on groundwater", 23-28 may 1994, Constanța, Romania, Bucharest University Press, 420-428.
- Popa Elena (1981) *Biostratigrafia formațiunilor mezozoice din partea orientala a Munților Pădurea Craiului*. Anuarul IGG LVIII, 203-282.
- Rusu, T. (1988) *Carstul din Munții Pădurea Craiului. Pe urmele apelor subterane*. Editura Dacia, Cluj-Napoca, 254 pages.